

A major existing shortfall of both models is the effective assumption of a zero incubation (or latent) period, i.e., once an individual is transferred from the susceptible to the infected class, that individual is assumed to be immediately capable of infecting other susceptibles. This is clearly unrealistic but could be overcome by the inclusion of a time delay factor.

If one assumes purely bubonic infection one should also attempt to model the rodent infective population and its interrelationship with the human population. However, such a compartment type model is likely to be inadequate due to the difficulty of obtaining realistic rodent data.

One should also perhaps consider the spatial aspects of plague infection. Certainly, for Eyam, the records clearly indicate that deaths occurred fairly rapidly among individual family units. The assumptions for the models presented here are based on perfect mixing which is clearly not accurate. Once again, however, one would have the difficulty of data availability in order to apply such models.

For present day outbreaks one would also need to include other factors such as vaccination control and variable population sizes. I am currently considering such models.

Readers requiring further background knowledge of mathematical epidemiology should refer to the papers by Bailey¹¹ and Anderson.¹² Both authors recognise the importance of modelling epidemic processes.

I am most indebted to Mr. K. Rennolls of the Forest Research Station, Farnham, Surrey, for discussions related to the setting up of the stochastic model described here.

References

1. Wood, W., "History of Eyam." Bell and Daldy, London, first published in 1865.
2. Daniel, C., "The Story of the Eyam Plague," published by the author, 1977.
3. Beaumont, R. M., "The Reverend William Mompesson 1638-1708, Hero of Eyam," Woods, Southwell, Notts., 1980.
4. Pollitzer, R., "Plague," World Health Organisation Monograph Series 22, 1954.
5. Raggett, G. F., *Applied Mathematical Modelling*, 1980, 4, 479-482.
6. Raggett, G. F., "A Mathematical Model of the Eyam Plague," Proceedings of 1st IASTED International Conference on Applied Modelling and Simulation, Lyon, France, Mesnard, G., Editor, 1981.
7. "Concise Medical Dictionary," Oxford Medical Publications, 1980.
8. Raggett, G. F., *Comp. Graphics*, 1979, 4, 149-154.
9. Bailey, N. T. J., "The Elements of Stochastic Processes with Applications to the Natural Sciences," Wiley, 1964.
10. Raggett, G. F., *Bulletin Appl. Stats.*, 1982, 9, No. 2, 212-225.
11. Bailey, N. T. J., "The Estimation of Parameters from Epidemic Models," in Bartlett, M. S., and Hiorns, R. W., Editors, "The Mathematical Theory of the Dynamics of Biological Populations I," Academic Press, London, 1973, pp. 253-267.
12. Anderson, R. M., "Infectious Disease Agents and Cyclic Fluctuations in Host Abundance," in Hiorns, R. W., and Cooke, D., Editors, "The Mathematical Theory of the Dynamics of Biological Populations II," Academic Press, London, 1981, pp. 47-80.

THE INSTITUTE OF MATHEMATICS AND ITS APPLICATIONS IRISH BRANCH



Dr. S.K.
Houston, FIMA



Mr. W. J.
Cromie, AFIMA

Chairman: Dr. S. K. Houston, FIMA (School of Mathematics, Ulster Polytechnic)
Secretary: Mr. W. J. Cromie, AFIMA (School of Mathematics, Ulster Polytechnic)

Branch activities

Throughout the year about eight meetings of the Branch are held covering a wide range of mathematical topics. The majority of the meetings are held in or near Belfast but some meetings are also held in Coleraine, Londonderry and Dublin. One meeting is arranged as a joint venture with the Royal Statistical Society. Occasional visits are also arranged and the most recent one was to the Planetarium and Observatory in Armagh. The Annual General Meeting of the Branch is usually held in May.

Coin Sequence Probabilities and Paradoxes

STANLEY COLLINGS, FIMA
The Open University, Milton Keynes




1. A COIN is tossed repeatedly, producing an endless random sequence of heads and tails (denoted hereafter by H and T). Players A and B wager on the first occurrence of their own nominated sub-sequence. Suppose A chooses HH and B chooses TH . A wins if the random sequence above produces H, H in consecutive places before it produces T, H in consecutive places in that order. As the probability of each sub-sequence is $\frac{1}{4}$, it might be thought that the wager was fair; but look at things this way.

A wins if the first two outcomes produce HH , and the probability of this is $\frac{1}{4}$.

B wins in all other cases; for as soon as any T occurs, TH must occur before any HH .

Therefore the odds in favour of B are 3:1.

2. Extending to sub-sequences of length 3, suppose A chooses TTT . B would then do better with any of HHT , HTH , THH , THT ; but HHT is best of all having odds of 7:1 in its favour. If we now look at every possibility for A , and at its best beater for B , we get the following figure (Fig. 1).



Thus we get an intransitive ring, but with "fly offs" producing something like a Catherine wheel. It follows that there is no best sub-sequence for A if B chooses second knowing A 's choice. But if B chooses completely at random, and A knows this is going to happen, he can maximise his own probability of success by selecting HHT or THH .

3. Before proceeding further, we should explain how to evaluate odds in general. These are done in terms of Conway Numbers (see reference 1) which are defined below. We shall work in terms of the particular example

$A-HTHH$
 $B-THTH$

The answers we require are expressed in terms of component figures denoted by AA , AB , BA , BB . To get the " AA " term, we write down A 's sub-sequence underneath itself getting

1
 $A-HTHH$
 $A-HTHH$

We put the digit 1 above the first H because the H leads a sequence which is identically the same as the four terms immediately below it. To get the next digit we

discard this H and slide the remaining three letters along to get

0
 THH
 $HTHH$

This time the digit is 0 because the T leads a sequence which is not the same as the three terms immediately below it.

Continuing with this process we get

0 1
 HH H
 $HTHH$ $HTHH$

Combining these results into a single display gives

1 0 0 1
 $HTHH$
 $HTHH$

We now interpret 1001 as a four digit number in the binary scale giving 9, and this is called the Conway Number corresponding to AA .

Doing this for all possible pairings of A and B gives

1 0 0 1 = 9	0 0 0 0 = 0
$A-HTHH$	$A-HTHH$
$A-HTHH$	$B-THTH$
1 0 1 0 = 10	0 1 0 1 = 5
$B-THTH$	$B-THTH$
$B-THTH$	$A-HTHH$

For convenience we shall write

$AA = 9$, $AB = 0$,
 $BB = 10$, $BA = 5$.

It is now the case (not mentioned in the Martin Gardner article¹) that the average waiting time to get A from scratch is 2. $AA = 18$. Also on average the further number of tosses required to complete or get an A having just completed a B is 2. $AA - 2$, $BA = 8$. It is however mentioned that the odds in favour of B winning the wager are $(AA - AB):(BB - BA)$.

Applying these results to our particular case,

A has an average waiting time of 18 to get $HTHH$,
 B has an average waiting time of 20 to get his $THTH$

implying an advantage to A . Yet the odds in favour of B 's wager are

$$\frac{9-0}{10-5} = \frac{9}{5}$$

By now, this is a paradox within a paradox.

4. Before proving these results ourselves, we shall generalise the situation. Tossing a penny is like rolling a 2-sided die. We shall consider what happens on rolling a k -sided die.

Theorem X

The expected waiting time to get a given sequence of integers from the set $\{1, 2, \dots, k\}$ is $k \cdot AA$, where AA stands for the corresponding Conway Number consisting entirely of 0's and 1's and interpreted in the scale of k .

Theorem Y

Given a sequence B to start with, the expected further number of rolls required to complete or produce the A sequence is $k \cdot AA - k \cdot BA$.

Proofs

We proceed by induction, supposing the results to be true for all pairs of sequences A, B of length $\leq n$. We then establish the results for length $n+1$.

(i) Let A, B both have length $n+1$. If they are identical, the A sequence is complete by virtue of B itself, so the further number of rolls required is zero. This fits in with theorem Y which says that the expected further number required is $k \cdot AA - k \cdot BA = k \cdot AA - k \cdot AA = 0$.

(ii) If they are not identical, BA must begin with a zero. Suppose the first r digits of BA are all zero, and that the $(r+1)$ th is 1. Then B consists of a "zeros-generating" sub-sequence C followed by a sub-sequence D generating a leading 1. Because of this, D is a leading sub-sequence of A .

(iii) By inspection

$$\begin{aligned} BA &= DA \\ &= DD \text{ as only the first } n+1-r \text{ terms of } A \\ &\text{are involved.} \end{aligned}$$

(iv) To get the sequence A , we must first get a D sub-sequence, and then follow it up successfully (though not necessarily on the first attempt). Therefore expected number of rolls to get A

$$\begin{aligned} &= \text{number required to get } D + \text{further number} \\ &\text{required to get from } D \text{ to } A \\ &= k \cdot DD + \text{further numbers required to get from} \\ &\text{ } D \text{ to } A \text{ (by the inductive hypothesis on} \\ &\text{ } \text{theorem X as } D \text{ has length } \leq n). \end{aligned}$$

$$\begin{aligned} \text{Therefore the number of rolls to get from } B \text{ to } A \\ &= \text{number to get from } D \text{ to } A, \text{ as } D \text{ is the only} \\ &\text{useful bit of } B \\ &= \text{number required to get } A \text{ from} \\ &\text{scratch} - k \cdot DD \\ &= \text{number required to get } A - k \cdot BA. \end{aligned} \quad (iii)$$

(v) The argument is essentially the same if B has length $\leq n$. The only difference occurs if BA has leading digit 1 making $D = B$; but it is still true that D has length $\leq n$ allowing us to apply the inductive hypothesis.

(vi) If A has length $\leq n$, B must have length $n+1$

$$\begin{aligned} \therefore BA &\text{ has leading digit } 0, \\ \therefore D &\text{ has length } \leq n \text{ and the earlier argument} \\ &\text{applies.} \end{aligned}$$

(vii) So far we have tacitly been assuming that $BA \neq 0$. If this is false, then there is no coincidence between any trailing sub-sequence of B and a leading sub-sequence of A . Although we have B to start with, it is of no assistance at all.

$$\begin{aligned} \therefore \text{number of rolls required to get from } B \text{ to } A \\ &= \text{number required to get } A \text{ from scratch} \\ &= \text{number required to get } A - k \cdot BA \text{ as} \\ &\quad BA = 0. \end{aligned}$$

(viii) We now temporarily break off the induction argument on theorem Y and apply it to theorem X. A has length $n+1$, and consists of a single outcome R followed by a sequence A' of length n . By inspection $AA = k^n + A'A$. The probability of getting the sequence A at any stage is $P(A) = 1/k^{n+1}$

$$\therefore \text{average distance between successive } A \text{ runs (as} \\ \text{measured between their respective last terms)} \\ \text{is } k^{n+1},$$

$$\begin{aligned} \therefore k^{n+1} &= \text{expected number of further rolls} \\ &\text{required to get from } A' \text{ to } A \\ &= \text{number to get } A - k \cdot A'A. \end{aligned} \quad (v)$$

$$\begin{aligned} \therefore \text{number of rolls required to get } A \text{ from scratch} \\ &= k^{n+1} + k \cdot A'A \\ &= k(k^n + A'A) = k \cdot AA. \end{aligned}$$

Thus the induction argument applies to theorem X.

(ix) Returning to theorem Y we already know that the number of rolls required to get from B to A = number required to get $A - k \cdot BA$. But this now equals $k \cdot AA - k \cdot BA$, and so the induction applies to theorem Y as well.

(x) Finally theorems X and Y are clearly true for $n = 1$; so they are true for all n .

5. We are now able to tackle the probabilities of the situation.

Theorem Z

The odds that the B sequence precedes the A sequence are given by $(AA - AB):(BB - BA)$.

Proof

Suppose the whole experiment is carried out N times, each time until both A and B appear; then if $P(B$ precedes $A) = q = 1 - p$, B will precede A in about Nq of the occasions. The average waiting time to get A is $k \cdot AA$ and to get B is $k \cdot BB$

$$\begin{aligned} \therefore \text{average distance between occurrences} \\ &= k(AA - BB), \text{ which can of course be positive or} \\ &\text{negative.} \end{aligned}$$

On those occasions when B precedes A , the average distance is the waiting time to get from B to A

$$= k \cdot AA - k \cdot BA.$$

When A precedes B , the average distance

$$\begin{aligned} &= -\text{waiting time to get from } A \text{ to } B \\ &= -(k \cdot BB - k \cdot AB). \end{aligned}$$

The weighted average of these last two must give the over-all average;

$$\begin{aligned} \therefore k(AA - BB) \\ &= q(k \cdot AA - k \cdot BA) - p(k \cdot BB - k \cdot AB). \end{aligned}$$

$$\therefore AA - BB = q \cdot AA - q \cdot BA - p \cdot BB + p \cdot AB,$$

$$\therefore p \cdot AA - q \cdot BB = p \cdot AB - q \cdot BA,$$

$$\therefore p(AA - AB) = q(BB - BA),$$

$$\therefore \text{odds in favour of } B \text{ preceding } A \text{ are}$$

$$q:p = (AA - AB):(BB - BA).$$

6. In theory, we can apply the above results to three person games. Thus suppose players A, B, C all choose their sequences, and that the probabilities of the various finishing orders are given by

$$P(ABC) = p_1, \quad P(BCA) = p_3, \quad P(CAB) = p_5$$

$$P(ACB) = p_2, \quad P(BAC) = p_4, \quad P(CBA) = p_6.$$

Then immediately we have

$$p_1 + p_2 + p_3 + p_4 + p_5 + p_6 = 1,$$

$$p_1 + p_2 + p_5 = P(A \text{ precedes } B),$$

$$p_3 + p_4 + p_6 = P(B \text{ precedes } A),$$

$$\therefore \frac{p_3 + p_4 + p_6}{p_1 + p_2 + p_5} = \text{odds on } B \text{ preceding } A$$

$$= \frac{AA - AB}{BB - BA}.$$

Similarly

$$\frac{p_2 + p_5 + p_6}{p_1 + p_3 + p_4} = \frac{BB - BC}{CC - CB}.$$

Again, in those games where A wins, the waiting time before B gets his sequence is $k(BB - AB)$ and before C gets his is $k(CC - AC)$;

$\therefore$ average distance = $k(BB - AB - CC + AC)$, the positive direction being from C to B .

But in these cases where A wins, the conditional odds on C beating B are $p_2:p_1$;

$\therefore$ by the same argument as in theorem Z,

$$\begin{aligned} k(BB - AB - CC + AC) \\ &= \frac{p_2 k}{p_1 + p_2} (BB - CB) - \frac{p_1 k}{p_1 + p_2} (CC - BC), \\ \therefore (p_1 + p_2)(BB - AB - CC + AC) \\ &= p_2(BB - CB) - p_1(CC - BC), \\ \therefore p_1(BB - BC - AB + AC) \\ &= p_2(CC - CB - AC + AB). \end{aligned}$$

With some licence, one could write this as

$$p_1(B - A)(B - C) = p_2(C - A)(C - B).$$

Similarly

$$\begin{aligned} p_3(C - B)(C - A) &= p_4(A - B)(A - C), \\ p_5(A - C)(A - B) &= p_6(B - C)(B - A). \end{aligned}$$

The multiplication between the brackets is of course non-commutative. We now have six equations for the 6 unknown probabilities p .

7. Returning to two players only, let us examine the expected time to the compound event $A \cup B$. It would be $k \cdot AA$ except for the fact that in a proportion q of the time, B precedes A thereby decreasing the above value. The longer A takes to occur on any particular occasion,

the greater is the probability that B will have preceded it; so the A and B events are dependent. Nevertheless in those cases where B does occur first, the outcomes subsequent to the arrival of B (and leading eventually to the occurrence of A) are completely random, and are independent of when the event B occurred. The average time from B to A is $k(AA - BA)$; hence in the proportion q of cases we overestimate the waiting time by the amount $k(AA - BA)$ if we persist with the value $k \cdot AA$. Therefore the required answer is

$$k \cdot AA - qk(AA - BA) = k(p \cdot AA + q \cdot BA);$$

equivalently it is $k(q \cdot BB + p \cdot AB)$. The fact that these two expressions are equal is equivalent to the statement of theorem Z.

8. By the same brand of argument, expected time to $A \cap B$ is

$$k \cdot AA + pk(BB - AB)$$

or

$$k \cdot BB + qk(AA - BA).$$

9. Another question is "what is the average time to get A given that it precedes B ." It is tempting to believe it is equal to the time to get $A \cup B$, and hence also to the average time to get B given that it precedes A . This would be very tidy, but unfortunately it is not true. If

$$A \text{ is } HH, \quad k = 2,$$

$$B \text{ is } TH, \quad p = \frac{1}{4}.$$

The average time to get A when it precedes B is 2 because A wins only when the first two outcomes are HH . The average time to get B when it precedes A is clearly greater than 2.

10. In a k -sided die situation, suppose that the two players can choose only between the sequences A and B . It is supposed that the sequence chosen by the first player already exists "on the table," and that Y pays to X in pennies the waiting time from here to his own choice. Thus if

$$k = 3,$$

$$A \text{ is } 213$$

$$B \text{ is } 121$$

then

$$100 = 9 \quad 000 = 0$$

$$A - 213 \quad A - 213$$

$$A - 213 \quad B - 121$$

$$101 = 10 \quad 010 = 3$$

$$B - 121 \quad B - 121$$

$$B - 121 \quad A - 213$$

and expectation pay-off matrix (from columns player to rows player) is

$$\begin{array}{cc|cc} & & \text{column choice} & & \\ & & A & B & \\ \text{row choice} & A & \begin{bmatrix} k(AA - AA) & k(BB - AB) \end{bmatrix} & = & \begin{bmatrix} 0 & 30 \\ 18 & 0 \end{bmatrix} \\ & B & \begin{bmatrix} k(AA - BA) & k(BB - BB) \end{bmatrix} & & \end{array}$$

Under standard mixed strategy game theory, the rows player should choose at random between first and second in the ratio 3:5.

Another way of analysing the game is for the rows player to argue:

"If column A is selected, I ought to get k . AA . The reason I do not is that my pre-sequence gives assistance which I grudge. If I choose A , my pay-off is $k(AA - AB)$ so my assistance amounts to k . AA . On the other hand with row B my assistance is only k . BA . I am hanged if I am going to assist my competitor any more than necessary. My strategy must therefore be to minimise from the pay-off matrix

$$\begin{bmatrix} AA & AB \\ BA & BB \end{bmatrix}$$

This is achieved by a random selection of rows with odds $(BB - BA):(AA - AB)$."

I think that such a player, one who does down his opponent even at the cost of his own personal gain, could be called spiteful.

Now let us change the scenario to where the row is selected not by human intellect but by blind chance. The obvious way of setting this up is to run a random sequence of rolls and to choose A when its precedes B , and B when it precedes A . By theorem Z this results in the odds $A:B = (BB - BA):(AA - AB)$. But these are precisely the odds to minimise rows assistance. Does this prove that blind chance is spiteful? Many of us have had occasions in our lives to suspect as much. Now we know!

Returning to our numerical example,

$$\begin{bmatrix} AA & AB \\ BA & BB \end{bmatrix} = \begin{bmatrix} 9 & 0 \\ 3 & 10 \end{bmatrix}$$

and the chance odds are

$$A:B = 7:9.$$

11. We have proved for a two person wager that the odds for precedence are the same as the spiteful odds in a mixed strategy game. Perhaps the equivalent argument can also go the other way. Let us test the situation in a simple numerical case with three players.

Take coin tossing where

player A chooses HH ,
player B chooses HT ,
player C chooses TT .

The equivalent pay-off matrix is

$$\begin{bmatrix} AA & AB & AC \\ BA & BB & BC \\ CA & CB & CC \end{bmatrix} = \begin{bmatrix} 3 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 3 \end{bmatrix}.$$

Odds must be selected for the rows to give the same return whichever column is chosen. Therefore

$$\begin{aligned} 3p &= p + 2q = q + 3r, \\ \therefore p &= q, \quad 2p &= 3r, \\ \therefore p:q:r &= 3:3:2. \end{aligned}$$

It is easily seen in this case that any variation in these odds allows "columns" to pay out less on average.

In (6), we had equations for the probabilities of the

various possible results with players A, B, C . Filling numerical values into those equations gives

$$\begin{aligned} p_1 + p_2 + p_3 + p_4 + p_5 + p_6 &= 1, \\ \frac{p_3 + p_4 + p_6}{p_1 + p_2 + p_5} &= \frac{AA - AB}{BB - BA} = \frac{2}{2} = 1, \\ \frac{p_2 + p_5 + p_6}{p_1 + p_3 + p_4} &= \frac{1}{3}, \end{aligned}$$

$$\begin{aligned} p_1(2 - 1 - 1 + 0) &= p_2(3 - 0 - 0 + 1), \\ p_3(3 - 0 - 1 + 0) &= p_4(3 - 0 - 0 + 1), \\ p_5(3 - 1 - 0 + 0) &= p_6(2 - 0 - 0 + 0). \end{aligned}$$

Simplifying,

$$\begin{aligned} p_1 + p_2 + p_3 + p_4 + p_5 + p_6 &= 1, \\ p_3 + p_4 + p_6 &= p_1 + p_2 + p_5 = \frac{1}{2}, \\ p_1 + p_3 + p_4 &= 3p_2 + 3p_5 + 3p_6, \\ p_2 &= 0, \\ p_3 &= 2p_4, \\ p_5 &= p_6, \end{aligned}$$

$$\therefore p_1 = \frac{3}{8}, \quad p_3 = \frac{2}{8}, \quad p_5 = \frac{1}{8}, \\ p_2 = 0, \quad p_4 = \frac{1}{8}, \quad p_6 = \frac{1}{8},$$

$$\begin{aligned} P(A \text{ wins}) &= p_1 + p_2 = \frac{3}{8}, \\ P(B \text{ wins}) &= p_3 + p_4 = \frac{3}{8}, \\ P(C \text{ wins}) &= p_5 + p_6 = \frac{2}{8}. \end{aligned}$$

giving exactly the same odds as above.

12. In (2) we considered each possible sub-sequence of length 3, located its best beater, and drew a diagram (Fig. 1) which contained an intransitive ring. Doing the same for length 4 gives (Fig. 2):




Fig. 2

Length 5 gives something even more complicated.

13. The above figure contains an intransitive hexagon. If we drop the condition about noting only best beaters, we can have intransitive polygons of other sizes, the largest is the 14-gon.

$$\begin{aligned} HHHT &\rightarrow THHT \rightarrow HHTT \rightarrow THTT \rightarrow HTHT \rightarrow \\ THTT &\rightarrow HHTT \rightarrow THTT \rightarrow HTTH \rightarrow TTHH \rightarrow \\ HHHT &\rightarrow THHT \rightarrow HHTT \rightarrow TTHH \rightarrow TTHH \rightarrow HHHT. \end{aligned}$$

The smallest are triangles; in fact we can get these from alternate vertices of the above hexagon

$$\begin{aligned} HHHT &\rightarrow HHTT \rightarrow TTHH (\rightarrow HHHT) \\ THTT &\rightarrow TTHH \rightarrow HHTT (\rightarrow THTT). \end{aligned}$$

14. It can be shown that the best choices for A against a purely random choice by B are $HHTT$ and $TTHH$. Next come the pair $HHTT$ and $TTHH$, and then the pair $HHHT$ and $TTHH$. It is interesting that these six are the vertices of the above intransitive hexagon. But while this list gives the preference order

$$\begin{aligned} HHTT \\ HHTT \\ HHHT \end{aligned}$$

it is apparent from the intransitive hexagon that on average $HHHT$ beats $HHTT$ and $HHTT$ beats $HHHT$ (which is the exactly reversed order!).

15. We turn now to the recent *Bulletin* article² on "Beetle." We shall adopt the same notation except that $W_{a,b}$ for us will denote the expected total number of rolls of an ordinary die to obtain a 5's and b 6's in any order (and not necessarily consecutively). In other words we proceed from 0, 0 to a, b ; also we work in terms of single rolls, not blocks of 6.

16. To get the compound event $5 \cup 6$, we expect to need 3 rolls. Therefore to get $2n$ of these results, we need $6n$ rolls. But however many rolls we in fact take, we shall be at the point

$$(0, 2n) \quad \text{with probability} \quad \binom{2n}{0} / 2^{2n}$$

$$(1, 2n-1) \quad \text{with probability} \quad \binom{2n}{1} / 2^{2n}$$

$$(2, 2n-2) \quad \text{with probability} \quad \binom{2n}{2} / 2^{2n}$$

$\vdots$

$$(n, n) \quad \text{with probability} \quad \binom{2n}{n} / 2^{2n}$$

$\vdots$

$$(2n, 0) \quad \text{with probability} \quad \binom{2n}{2n} / 2^{2n}$$

$$\begin{aligned} \therefore W_{n,n} &= 6n + 6n \binom{2n}{0} / 4^n \\ &\quad + 6(n-1) \binom{2n}{1} / 4^n + \dots + 6 \binom{2n}{n-1} / 4^n \\ &\quad + 0 \cdot \binom{2n}{n} / 4^n + 6 \binom{2n}{n+1} / 4^n \\ &\quad + \dots + 6n \binom{2n}{2n} / 4^n \end{aligned}$$

because from

$$\begin{aligned} (0, 2n) &\text{ we still need } n \text{ 5's} \\ (1, 2n-1) &\text{ we still need } (n-1) \text{ 5's} \\ &\vdots \\ (n, n) &\text{ we need nothing more} \\ &\vdots \\ (2n, 0) &\text{ we still need } n \text{ 6's.} \end{aligned}$$

$$\begin{aligned} \therefore W_{n,n} &= 6n + \frac{2 \cdot 6}{4^n} \left\{ n \binom{2n}{0} + (n-1) \binom{2n}{1} \right. \\ &\quad \left. + \dots + 1 \binom{2n}{n-1} \right\} \\ &= 6n + \frac{12}{4^n} K_n. \end{aligned}$$

where

$$\begin{aligned} K_n &= n \left\{ \binom{2n}{0} + \binom{2n}{1} + \dots + \binom{2n}{n} \right\} \\ &\quad - \left\{ 0 \cdot \binom{2n}{0} + 1 \binom{2n}{1} + \dots + n \binom{2n}{n} \right\} \\ &= n \left\{ \frac{1}{2} \cdot 2^{2n} + \frac{1}{2} \binom{2n}{n} \right\} \\ &\quad - \left\{ \frac{(2n)!}{0!(2n-1)!} + \frac{(2n)!}{1!(2n-2)!} + \dots + \frac{(2n)!}{(n-1)!n!} \right\} \\ &= \frac{n}{2} \left\{ 4^n + \binom{2n}{n} \right\} - 2n \left\{ \binom{2n-1}{0} \right. \\ &\quad \left. + \binom{2n-1}{1} + \dots + \binom{2n-1}{n-1} \right\} \\ &= \frac{n}{2} \left\{ 4^n + \binom{2n}{n} \right\} - 2n \left\{ \frac{1}{2} \cdot 2^{2n-1} \right\} \\ \therefore W_{n,n} &= 6n + \frac{12}{4^n} \cdot \frac{n}{2} \left\{ 4^n + \binom{2n}{n} \right\} - \frac{12}{4^n} \cdot n \cdot 2^{2n-1} \\ &= 6n + 6n + 6n \binom{2n}{n} / 4^n - 6n \\ &= 6n + 6n \binom{2n}{n} / 4^n. \end{aligned}$$

$$\begin{aligned} 17. \quad W_{n,n} &= 3 + \frac{1}{2} W_{n-1,n} + \frac{1}{2} W_{n,n-1} \\ &= 3 + W_{n-1,n} \quad \text{by symmetry,} \\ \therefore W_{n-1,n} &= W_{n,n} - 3. \end{aligned}$$

18. We know that

$$\begin{aligned} W_{n,1} &= 6n + 6/2^n, \\ W_{n,2} &= 6n + 6(n+4)/2^{n+1}. \end{aligned}$$

Suppose inductively that

$$W_{n,3} = 6n + 3(n^2 + 9n + 24)/2^{n+2}.$$

This is true for $n = 1$, as then

$$\begin{aligned} W_{1,3} &= 6 + 3(1 + 9 + 24)/2^3 \\ &= 6 + \frac{3 \cdot 34}{8} = 18\frac{3}{4} = W_{3,1}. \end{aligned}$$

Also,

$$\begin{aligned} W_{n+1,3} &= 3 + \frac{1}{2} W_{n,3} + \frac{1}{2} W_{n+1,2} \\ &= 3 + 3n + 3(n^2 + 9n + 24)/2^{n+3} \\ &\quad + 3(n+1) + 6(n+5)/2^{n+3} \\ &= 6(n+1) + 3\{n^2 + 9n + 24 + 2(n+5)\}/2^{n+3} \\ &= 6(n+1) + 3\{(n+1)^2 + 9(n+1) + 24\}/2^{n+3} \end{aligned}$$

which completes the induction argument.

Again, suppose inductively that

$$W_{n,4} = 6n + (n^3 + 15n^2 + 86n + 192)/2^{n+3}.$$

This is true for $n = 1$, as then

$$W_{1,4} = 6 + 294/2^4 = 24\frac{3}{8} = W_{4,1}.$$

Also

$$\begin{aligned} W_{n+1,4} &= 3 + \frac{1}{2} W_{n,4} + \frac{1}{2} W_{n+1,3} \\ &= 3 + 3n + (n^3 + 15n^2 + 86n + 192)/2^{n+4} + 3(n+1) \end{aligned}$$

$$\begin{aligned}
& + 3(n^2 + 11n + 34)/2^{n+4} \\
& = 6(n+1) + (n^3 + 18n^2 + 119n + 294)/2^{n+4} \\
& = 6(n+1) + \{(n+1)^3 + 15(n+1)^2 \\
& \quad + 86(n+1) + 192\}/2^{n+4}
\end{aligned}$$

which completes the induction argument. It is obvious from this that $W_{n,r}$ contains a polynomial in n of degree $r-1$, but it is not obvious what that polynomial is.

19. In 4(iii) of the paper, the stated waiting time of $n^2 + 1$

for 3 consecutive wins out of n equally competent teams is surely a misprint for $n^2 + n + 1$.

20. From the same sub-section, a closer upper bound for the waiting time for any 3 wins is $1 + 2n$. For $n = 2$ this gives 5, whereas the exact answer is $4\frac{1}{8}$. For $n = 3$, the answer is 5.05 (E. and O.E.).

References

1. Gardner, M., *Scientific American*, October 1974, pp. 120-125.
2. Watson, F. R., *Bull. IMA*, 1982, 18, 18-25.

THE INSTITUTE OF MATHEMATICS AND ITS APPLICATIONS

IMA JOURNAL OF NUMERICAL ANALYSIS

Editors: Professor K. W. Morton, FIMA (University of Reading) and Professor M. J. D. Powell, FIMA (University of Cambridge)

The following papers will be published in the 1982 issues of the *IMA Journal of Numerical Analysis*.

- "A Formula Approximating the Root of a Function"—J. E. Dawson (CSIRO, Sydney, Australia)
- "Further Developments of RP and AP Error Analysis"—F. W. J. Olver (The University of Maryland and the National Bureau of Standards)
- "A Mixed Method for the Elastoplastic Torsion Problem"—A. Bermudez (Universidad de Santiago de Compostela, Spain)
- "Geometrically Derived Difference Formulae for the Numerical Integration of Trajectory Problems"—R. J. Y. McLeod (New Mexico State University, USA) and J. M. Sanz-Serna (Universidad del Pais Vasco Lejona, Spain)
- "A Special Class of Multistep Runge-Kutta Methods with Extended Real Stability Interval"—P. J. van der Houwen and B. P. Sommeijer (Stichting Mathematisch Centrum, Amsterdam, The Netherlands)
- "A Numerical Method for Conformal Mapping"—N. V. Challis (Sheffield City Polytechnic) and D. M. Burley (University of Sheffield)
- "The Construction of Reducible Quadrature Rules for Volterra Integral and Integro-differential Equations"—P. H. M. Wolkenfelt (Stichting Mathematisch Centrum, Amsterdam, The Netherlands)
- "Mono-Implicit Runge-Kutta Formulae for the Numerical Integration of Stiff Differential Systems"—J. R. Cash and A. Singhal (Imperial College, London)
- "A Complementary Solution to the Dam Problem"—R. W. Thatcher and S. L. Askew (The University of Manchester Institute of Science and Technology, Manchester)
- "Piecewise Rational Quadratic Interpolation to Monotonic Data"—J. A. Gregory (Brunel University, Uxbridge) and R. Delbourgo (Middlesex Polytechnic, London)
- "Rational Approximations from the d-Transformation"—A. Sidi (Israel Institute of Technology) and D. Levin (Tel Aviv University, Israel)
- "The Use of Auto-correlation for Pseudo-rank Determination in Noisy Ill-conditioned Linear Least-squares Problems"—M. L. Baart (National Research Institute for Mathematical Sciences, South Africa)

The 1982 (Volume 2, nos. 1-4) issues of the *IMA Journal of Numerical Analysis* can be obtained from the IMA at price £22 (members); £53.60 (non-members).

TRANSFER TO ASSOCIATE FELLOWSHIP

Graduate members are reminded of the regulations relating to Associate Fellowship. All Graduate members are deemed to comply with the academic requirements for Associate Fellowship and their eligibility for transfer is, therefore, dependent wholly on their professional experience. The minimum period of responsible experience is three years. All applications individually considered and the Secretary will always be glad to advise. Forms of application can be obtained from the Secretary.

Bharati Krishna's System of Vedic Mathematics

J. H. PICKLES, FIMA

Central Electricity Generating Board, Leatherhead

A. P. NICHOLAS

Polytechnic of North London

and

K. R. WILLIAMS

Department of Physics and Astronomy, University College London

1. Introduction

"THE nine Indian figures are 9 8 7 6 5 4 3 2 1. With these nine figures and the sign 0, ..., any number may be written as is demonstrated below." So wrote Leonardo Fibonacci in his "*Liber Abaci*," published in 1202, which introduced the decimal numeral system into Europe. This advance in notation was vital to the subsequent progress in mathematical thought and technique.

What is perhaps surprising, as pointed out by the Indian scholar and teacher Bharati Krishna (1884-1960), is that the methods we use for the simple operations of arithmetic—multiplication, division, and so on—are still not those best suited to use with the decimal system. In his book "*Vedic Mathematics*" Bharati Krishna¹ develops, apparently from the original Indian tradition, a range of attractive alternative techniques. These substantially reduce the computational drudgery associated with mental or pencil-and-paper arithmetic and seem to provide an effective antidote to the phenomenon of "overdependence on the calculator for trivial calculations."²

2. Examples

Multiplication

One very simple method applies particularly to those parts of the multiplication tables which seem less than trivial to a fair proportion of the school population.³ Consider the multiplication of 9×7 . The procedure is to subtract both these digits from the decimal base 10, giving deficiencies of 1 and 3, which are written out thus

$$\begin{array}{r}
9-1 \\
7-3 \\
\hline
/
\end{array}$$

in anticipation of a two-digit answer. To find the first digit, "cross-subtract" the 3 from the 9 or (to take the easier option!) the 1 from the 7: $9-3 = 7-1 = 6$. For the second digit, just multiply the two deficiencies:

$$\begin{array}{r}
9 \quad 1 \\
\times \\
7 \quad 3 \\
\hline
6 \quad 3
\end{array}$$

In terms of algebra,

$$\begin{aligned}
(x-1)(x-3) &= x(x-1-3) + 1 \times 3 \\
&= 6x + 3,
\end{aligned}$$



where x is the decimal base 10. From the "Vedic" point of view, the answer is not so much the single magnitude 63 as the combination of algebraic coefficients, 6 and 3. The superficially irreducible calculation 9×7 breaks down into what, in practice, are simpler steps for the calculation of the separate coefficients.

Essentially the same principle applies in relation to bases of 100, 1000, ..., etc. or to "working" bases which are fractions or multiples thereof. For example, to multiply 963×989

$$\begin{array}{r}
963-037 \\
989-011 \\
\hline
952/407
\end{array}$$

The deficiencies from the base 1000 are here calculated, digit by digit, or coefficient by coefficient, according to the rule or "sutra": "all from nine and the last from ten." Thus $9-9 = 0$, $9-6 = 3$, $10-3 = 7$, etc. Again, with a working base of 20, 19×17 becomes

$$\begin{array}{r}
19-1 \\
17-3 \\
\hline
16/3
\end{array}$$

and doubling the left hand digit pair gives the answer 32/3.

The scope of this device can be stretched further to cater, for example, for positive deficiencies, or for the "carrying" of surplus digits, but there are still cases where no very convenient working base can be found. Here the Vedic mathematician turns to a "sutra" which translates as "vertically and cross-wise." Take 13×23 :

$$\begin{array}{r}
1 \quad 3 \\
2 \quad 3 \\
\hline
\end{array}$$

The sutra finds the digits of the answer by following the pattern of the coefficients in the algebraic multiplication $(x+3)(2x+3)$. Thus

- (i) for the units digit, $3 \times 3 = 9$ (i.e., vertically ↓)
- (ii) for the tens digit, $(1 \times 3) + (2 \times 3) = 9$ (i.e., crosswise ✕)
- (iii) for the hundreds digit, $1 \times 2 = 2$ (vertically again ↑)

giving

$$\begin{array}{r}
1 \quad 3 \\
2 \quad 3 \\
\hline
299
\end{array}$$